Black, S. H., and D. M. Vaughan. 2005. Species Profile: *Icaricia icarioides fenderi*. In Shepherd, M. D., D. M. Vaughan, and S. H. Black (Eds). *Red List of Pollinator Insects of North America*. CD-ROM Version 1 (May 2005). Portland, OR: The Xerces Society for Invertebrate Conservation.

Icaricia icarioides fenderi Macy, 1931 Fender's Blue (Lycaenidae: Polyommatinae: Polyommatini)



Profile prepared by Scott Hoffman Black and Mace Vaughan The Xerces Society for Invertebrate Conservation

SUMMARY

Once thought to be extinct, the Fender's Blue but terfly was known only from collections made between 1929 and 1937, until it was rediscovered in 1989 by Paul Hammond. Fender's Blue is a butterfly that formerly thrived in the prairies of Oregon's Willamette Valley, but during the past 140 years an estimated 99 percent of this native prairie has been turned into farmland or otherwise developed.

Habitat loss from agriculture and urban development poses the greatest threat to the Fender's Blue. Another major threat to the butterfly is the invasion of exotic plants. Invasive species, such as Himalayan blackberry and Scotch broom, have been out-competing and displacing the Kincaid's lupine and other native wildflowers in upland prairies where the butterflies live.

CONSERVATION STATUS Xerces Red List Status: Critically Imperiled Other Rankings:

| Canada – Species at Risk Act: | N/A |
|-------------------------------|------------|
| Canada – provincial status: | N/A |
| Mexico: | N/A |
| USA – Endangered Species Act: | Endangered |
| USA – state status: | N/A |
| NatureServe: | G5T1 |
| IUCN Red List: | N/A |
| | |

SPECIES PROFILE

DESCRIPTION

The Fender's Blue is a small butterfly in the family Lycaenidae (gossamer-wings), with a wingspan of about one inch. The upperside of the wings of male Fender's Blue butterflies are predominately iridescent sky blue. Females have rusty brown wings. The dorsal surface of both sexes have a black border within a white fringe and the underside is pearly gray to dirty chalk, with black and brown spots, outlined in white.

TAXONOMIC STATUS

Icaricia icarioides fenderi Macy, 1931.

LIFE HISTORY

Fender's Blue lives primarily in the upland prairies of Oregon's Willamette Valley. The Kincaid's lupine (*Lupinus oreganus* var. *kincaidii*)—a wildflower formally listed as threatened because of extensive habitat loss—is the primary larval hostplant and is critical to the survival of this butterfly. Caterpillars have also been found on sicklekeeled lupine (*Lupinus albicaulis*) and the longspurred lupine (*Lupinus arbustus*).

Adult butterflies begin laying their eggs, primarily on the Kincaid's lupine, during the month of May. Once the larvae hatch, the caterpillars feed briefly until the plants begin to senesce in early July. At that time, they drop to the base of the plants and enter diapause (a stage in the butterfly's life when its growth and development are suspended) for the fall and winter. In February or March the larvae come out of diapause and continue their development. The larvae then enter their pupal stage and subsequently emerge as adult butterflies in May. A Fender's Blue butterfly completes its life cycle in one year.

DISTRIBUTION

The Fender's Blue is endemic to the native upland prairies of the Willamette Valley in Oregon. Only one-tenth of one percent of the original habitat once available to the Fender's Blue and Kincaid's lupine, or approximately 400 hectares of prairie, exists today.

THREATS

The principal threats are loss of habitat and degradation of the vegetation communities on habitat that remains.

CONSERVATION STATUS

Fender's Blue occurs in 32 small sites totaling 408 acres (approximately 165 hectares). Twenty-four of these populations occupy sites of eight acres or less. Most of these populations are in decline, and 19 contain fewer than 50 individuals. Fourteen sites are on federal, state, county, or city lands; the remainder are privately owned.

In surveys in 2001 many populations showed declines that are believed to be the result of the previous winter's drought conditions. Benton County had four stable, three decreasing, and one increasing population. Lane County had one stable, three decreasing, and two increasing populations. Active management—including mowing and burning to remove competitive weeds—is occurring at several sites.

Fender's Blue is federally listed as endangered. A recovery plan for Fender's Blue, Kincaid's lupine, and Willamette daisy is currently under development by Fish and Wildlife Service.

Recovery Plan (ESA): In development. Critical Habitat (ESA): None designated

Oregon's Threatened and Endangered List statute excludes insects from listing, so Fender's Blue is not provided with any protection at the state level. (Kincaid's lupine, however, is listed in Oregon as threatened.)

CONSERVATION NEEDS

Efforts to restore degraded prairie habitat and to remove non-native plants will be critical to bringing this species closer to recovery.

RESEARCH NEEDS

Currently, scientists are engaged in an active research program to study the habitat needs of this butterfly's hostplant. In addition, scientists with the Nature Conservancy, with the help of Xerces Society volunteers, are looking for new populations.

Scientific research into breeding the Fender's Blue butterfly are in their initial stage. Cheryl Schultz, Assistant Professor of Zoology, University of Washington, is heading up this effort.

RESOURCES

CONTACTS

U.S. Fish and Wildlife Service: Amy Horstman, Oregon Fish and Wildlife Office, Telephone: (503) 231-6179.

The Nature Conservancy of Oregon: Cathy MacDonald, Portland, OR. (503) 230-1221.

Washington State University: Cheryl Schultz, PhD, Assistant Professor, Zoology, WSU Vancouver, WA 98686-9600. Telephone: 360-546-9525; email: schultzc@vancouver.wsu.edu.

Willamette Resources and Educational Network: Pat Johnston, Project Manager, Bureau of Land Management, West Eugene Wetlands Project Office, 751 S. Danebo, Eugene, OR 97330. Telephone: 541-683-6181; email: pat_Johnston@or.blm.gov.

REFERENCES

Black, S. H., and M. Vaughan. 2004. Protecting the Northwest prairies and their butterflies. *Wings* 27(1):18-22.

Hammond, P. C. 1995. *1994 Study of the Fender's Blue Butterfly* (Icaricia icarioides fenderi) *in Benton, Polk, and Yamhill Counties*. Report to the Oregon Natural Heritage Program and the U.S. Fish and Wildlife Service. 34 pages.

Hammond, P. C., and M. V. Wilson. 1992. Biodiversity and the conservation of endangered species. *Northwest Environmental Journal* 8:174-176.

Hammond, P. C., and M. V. Wilson. 1993. *Status of the Fender's Blue Butterfly*. Report to the U.S. Fish and Wildlife Service. 66 pages.

Hitchcock, C. L., and A. Cronquist. 1973. *Flora of the Pacific Northwest*. University of Washington Press, Seattle. 730 pages.

Oregon Natural Heritage Program. 1995. *Rare, Threatened and Endangered Plants and Animals of Oregon*. Oregon Natural Heritage Program, Portland, Oregon. 84 pages.

Schultz, C. B. 1994. *Status of the Fender's Blue Butterfly* (Icaricia icarioides fenderi) *in Eugene, Oregon: a Species at Risk*. Report to the Oregon Natural Heritage Program and the U.S. Fish and Wildlife Service. 43 pages.

Schultz, C. B. 1995. *Status of the Fender's Blue Butterfly* (Icaricia icarioides fenderi) *in Lane County, Oregon: a Year of Declines*. Report to the Oregon Natural Heritage Program and the U.S. Fish and Wildlife Service. 58 pages.

Wilson, M. V., P. C. Hammond, and J. B. Kauffman. 1992. The value of restoration and management in protecting biodiversity. *Northwest Environmental Journal* 8:201-202.

WEBSITES

U.S. Fish & Wildlife Service; Threatened and Endangered Species System: Fender's Blue Butterfly.

http://ecos.fws.gov/SpeciesProfile?spcode=I0IS

(Accessed 3/28/05)

NatureServe Explorer. <u>http://www.natureserve.org/explorer/</u> (Accessed 3/28/05)

Oregon State University; Perspectives: Fender's Blue Butterfly. http://oregonstate.edu/~wilsomar/Persp_FBB.htm (Accessed 11/2/2004)

Pacific Biodiversity Institute; Endangered Species Information Network: Fender's Blue Butterfly. <u>http://www.pacificbio.org/ESIN/Butterflies/Fender-Blue/Fender_Website_allframes/fender.htm</u> (Accessed 3/28/05)

Media articles University of California, Santa Barbara; "Endangered Butterfly Needs Special Environment." <u>http://www.instadv.ucsb.edu/pa/display.aspx?pkey=698</u> (Accessed 3/28/05)

CNN.com; "Double Jeopardy: Endangered Butterfly Depends On Endangered Plants." <u>http://www.cnn.com/2000/NATURE/06/26/butterflies.enn/</u> (Accessed 11/2/2004)

The Register-Guard, Eugene, OR; "Rare butterfly thrives in preserve," by Lance Robertson. <u>http://www.4j.lane.edu/partners/eweb/ttr/butterflies/butterfly.html</u> (Accessed 11/2/2004)